

Erratalista "Studies of Risks Associated with Atrial Fibrillation"

Sida 15. Står: (that is, mortality, strokes and hospitalisation)

Skall stå: (other than quality of life, such as mortality, strokes and hospitalisations)

Sida 15. Står: Limited data on hard endpoints, other than quality of life, such as mortality and stroke, and long-term freedom of AF limits its use.

Skall stå: Limited availability, limited data on hard endpoints such as mortality and stroke and long-term freedom of AF limits its use.

Sida 19. Statistical analysis paper I and II

Lägg till: Means and proportions for continuous and categorical variables were calculated. The above figures are presented for each period of AF admission, gender and comorbidity. The association of each period of AF admission with 1-year mortality, independently of age, gender and comorbidity are presented as hazard ratios with first period as reference and estimated through a Cox regression model. The same model implies also the independent association of the co-morbidity variables with 1-year mortality.

Sida 23. Står: "hyperthyroidism in only 0.4%"

Skall stå: "hyperthyroidism in only 1.0%"

Sida 30. Table 10. Står: *Includes chronic obstructive pulmonary disease and asthma

Skall stå: 1. Myocardial infarction and unstable angina pectoris. 2. chronic obstructive pulmonary disease.

Sida 31 Table 11. Båda kolumner för n events har försvunnit. Var god se delarbete II för den korrekta tabellen.

Sida 43. Står "is associated high but varying prevalence"

Skall stå: "is associated with high but varying prevalence"

Sida 58 Reference 75. Författare: Stroke Risk in Atrial Fibrillation Working Group

Sida 63 Reference 134. Författare: Meta-analysis Research Group in Echocardiography (MeRGE) Heart Failure Collaborators

Arbete 1.

Sida 11. Reference 28. Författare: Stroke Risk in Atrial Fibrillation Working Group

Arbete 2.

Sida 3. Table 2. Ersätt * and ** in tabellfoten med 1 and 2