

Contact allergy to propolis and beeswax. Diagnostics, occurrence, and chemistry.

Akademisk avhandling

Som för avläggande av medicine doktorsexamen vid Sahlgrenska akademien, Göteborgs universitet kommer att offentligens försvaras i Hörsal Arvid Carlsson, Medicinaregatan 3, den 21/10 2022, klockan 13.00

av Gunnar Nyman

Fakultetsopponent:

Åke Svensson, Docent

Lunds universitet, Lund, Sverige

Avhandlingen baseras på följande delarbeten

- I. G NYMAN, M TANG, A INEROT, A OSMANCEVIC, P MALMBERG, L HAGVALL, 2019, Contact allergy to beeswax and propolis among patients with cheilitis or facial dermatitis. *Contact Dermatitis* 2019; 81:110-116
- II. G NYMAN, S OLDBERG-WAGNER, K PRYSTUPA-CHALKIDIS, K RYBERG, L HAGVALL, Contact Allergy in Western Sweden to Propolis of Four Different Origins *Acta Dermato-Venereologica* 2020; 100: adv00256
- III. G NYMAN, AM GIMÉNEZ-ARNAU, J GRIGAITIENĖ, L MALINAUSKIENE, E PAULSEN and L HAGVALL, 2021, Patch Testing with Propolis of Different Geographical Origins in a Baseline Series. *Acta Dermato-Venereologica* 2021; 101: adv00591

**SAHLGRENKA AKADEMIN
INSTITUTIONEN FÖR KLINISKA VETENSKAPER**



Contact allergy to propolis and beeswax. Diagnostics, occurrence, and chemistry.

Gunnar Nyman

Avdelningen för dermatologi och venerologi, Institutionen för kliniska vetenskaper, Sahlgrenska akademien, Göteborgs universitet, Sverige, 2022.

Abstract

Cosmetics, including biocosmetics and “natural” skin remedies are common causes of contact dermatitis. Two frequent ingredients in biocosmetics are beeswax and propolis, which are known causes of contact allergy. These ingredients share components, with each other, and with plant-derived products and fragrances. The composition of propolis varies according to the plants growing in the vicinity of the beehive. How this variation influences the tendency of the propolis to cause contact allergy is not known.

The aim of this dissertation was to investigate contact allergy to beeswax and to propolis of different geographical origins. The method for demonstrating contact allergy is patch testing, a standardized provocation test.

We studied the frequency of contact allergy to beeswax and propolis in patients with cheilitis and facial dermatitis. Two studies regarding contact allergy to propolis of different geographical origins (from China, Lithuania, North America, and Sweden) were carried out. One included patients referred for patch testing in West Sweden, and the other included such patients in Denmark, Lithuania, and Spain. High frequencies of contact allergy to beeswax and propolis, were found. Small differences between the four types of propolis were demonstrated in each country. Propolis originating from China, and from Lithuania gave statistically higher frequencies of contact allergy than propolis from Sweden in the four countries together. High frequencies of concomitant positive patch test reactions between propolis and plant-related products and fragrances were found.

Beeswax and propolis should be considered as causes of contact allergy in patients with cheilitis and facial dermatitis. Patch testing with just one propolis preparation detected about half of the propolis-allergic patients. Contact allergy to propolis was so common that it should be considered for inclusion in regular patch testing.

Keywords: Beeswax, caffeic acid, caffeic acid 1,1-dimethylallyl ester, Caffeic acid phenylethyl ester, cera alba, cera flava, cheilitis, concomitant reactivity, patch test, propolis.